

A Community-Based Study of the Rate of Beta-Hemolytic Group A Streptococcal Infections in Symptomatic and Asymptomatic School Children

Sir,

Group A streptococcus (GAS), *Streptococcus pyogenes*, is a human pathogen that known to cause a wide spectrum of infections. Disease spectrum ranges from relatively benign infections to severe invasive diseases leading to serious post streptococcus sequelae, like acute rheumatic fever and acute glomerulonephritis and rheumatic fever (RF), rheumatic heart disease (RHD).^[1] GAS are the most frequently isolated pathogens in acute pharyngotonsillitis in school-aged children,^[2] and streptococcal pharyngitis is one of the most common childhood bacterial infections.

A surveillance study of beta-hemolytic streptococci (BHS), infections was conducted in school children aged 5-15 years from two different government public schools in Arasur village, located near the city of Gulbarga in Karnataka, South India. The approximate distance between Gulbarga and Arasur village is approximately 30 km. Arasur village has poor hygienic conditions and a lack of basic treatment facilities situated in a typical agricultural farmland represent

spring climate condition with temperature varying 22-27°C around the year. A total of 1052 children were randomly selected, including 997 asymptomatic children (normal) and 55 with symptoms of pharyngitis (symptomatic) during entire study period from August 2009 to January 2010. The study was designed and approved by ethical committee (institutional review board) of our institution. Throat swabs and blood samples (~3 mL) were collected from each child who presented with clinical signs or symptoms of sore throat, fever, chills, malaise, erythema, and swelling of the pharyngeal mucosa were included. Samples were also collected from asymptomatic (normal) children who were followed for couple of weeks to look into any possibility of developing pharyngitis.

Isolates were confirmed by the Streptococcus Laboratory, Department of Clinical Microbiology, Christian Medical College, Vellore, Tamil Nadu, using standardized methods.^[3]

Anti-streptolysin O (ASO) antibodies were detected by commercial test kits according to manufacturer's instructions (Tulip Diagnostics, Goa, India). Among the asymptomatic children, BHS were isolated from 113 (11.3%) of the 997 children screened. Of these isolated BHS, 35 (30.9%) belonged to group A, 11 (9.7%) to group B, 7 (6.1%) to group F, 21 (18.5%) to group C, and 39 (34.5%) to group G. A total of 46 BHS were isolates from 55 symptomatic children. Of these BHS, 18 (39.1%) belonged to group A, 6 (13%) to group C,

Table 1a: Rate of Lance field beta-hemolytic streptococci among school children

School children	Symptomatic carriers	Asymptomatic carriers	Total
No. screened	55	997	1052
BHS	46 (83.6%)	113 (11.3%)	159
GAS	18 (39.1%)	35 (30.9%)	53
GBS	0	11 (9.7%)	11
GCS	6 (13.0%)	21 (18.5%)	27
GFS	0	7 (6.1%)	7
GGG	22 (47.8%)	39 (34.5%)	61

BHS: Beta hemolytic Streptococci, GAS: Group A streptococcus, GBS: Group B streptococcus, GCS: Group C streptococcus, GFS: Group F streptococcus, GGS: Group A, B, C, F and G Streptococci

Table 1b: ASO positivity among children with GAS and non-GAS throat infection.

ASO	Symptomatic (n=46)		Asymptomatic (n=113)	
	GAS+	Non GAS+	GAS+	Non GAS+
Positive	8	0	4	0
Negative	10	28	31	78
Total	18	28	35	78

ASO: Antistreptolysin O, GAS: Group A streptococci, ASO+and GAS positivity among symptomatic children (8/18) and asymptomatic children (4/35), was statistically significant, with $P=0.0065 (<0.05)$.

and 22 (47.8%) to group G [Table 1a]. None of the BHS isolated from symptomatic school children belonged to group B or group F. Therefore, in this study, the overall incidence rates of GAS, group C streptococci (GCS), and group G streptococci (GGS) infection among symptomatic and asymptomatic children were found to be 5%, 2.5%, and 5.7%, respectively [Table 1b]. Eight (17.3%) of the 46 symptomatic children were ASO positive; (with titers of 200 IU/mL and 2 with titers of 400 IU/mL) and all eight children were also positive for GAS. Of the 113 asymptomatic children, four (3.5%) were positive for ASO. Their titers were 200 IU/mL ($n = 2$), 600 IU/mL, and 800 IU/mL. All four children were also positive for GAS. None of the children with non-GAS demonstrated ASO positivity. Insufficient hygienic facilities and lack of other basic amenities at schools and children from lower socioeconomic classes might be the reason for high GAS carriage in current study. GGS/GCS are generally considered to be commensals; however, due to horizontal transfer of genetic material with GAS (Davies *et al.*, 2005)^[4] GGS and GCS share many virulence factors same tissue niche in humans, and cause a similar spectrum of diseases.

In the final analysis, a laboratory diagnosis of GAS pharyngitis can only be made by recovering the organism in high numbers (>100 colonies on sheep blood agar) in the presence of clinical signs and symptoms. However in endemic regions, this may also be due to a viral infection with heavy GAS colonization. If ASO testing is carried out in such cases, it can act as a useful diagnostic adjunct for accurately confirming the cause of the infection.^[5] We conclude that regular surveillance of BHS throat infections in rural school children may be helpful in designing effective preventive strategies for poststreptococcal sequelae.

**Lakshmana Gowda Krishnappa^{1,2},
Mohammed Ali M Marie¹, James John³,
Shivannavar Channappa Thippa²,
Sangeetha Gopalkrishnan⁴,
Brahmadathan Kootallur Narayan³**

¹Departments of Clinical Lab Sciences, College of Applied Medical Sciences, King Saud University, Riyadh, Kingdom of Saudi Arabia, ³Clinical Microbiology, Christian Medical College and Hospital, Vellore, Tamil Nadu, ²Microbiology, Gulbarga University, Gulbarga, Karnataka, ⁴Microbiology, Central Leprosy Training and Research Institute, Chennai, Tamil Nadu, India

Address for correspondence:

James John, Ph.D

E-mail: james_cmc2002@yahoo.co.in

REFERENCES

1. Steer AC, Danchin MH, Carapetis JR. Group A streptococcal infections in children. *J Paediatr Child Health* 2007;43:203-13.
2. Pandian R, Brahmadathan KN, Mathai E. Throat and Nose. In: 13. Kang HP, Dunne WM, editors. *Stability of Repetitive-Sequence PCR Myer's and Koshi's manual of Diagnostic Procedures in Medical Microbiology and Immunology/Serology*. Faculty, Department of Clinical Microbiology, Christian Medical College, Vellore, India. Pondicherry: All India Press; 2001. p. 52-6.
3. Johnson DR, Kaplan EL, Sramek J, Bicova R, Havlicek J, Havlikova, *et al.* Laboratory diagnosis of group A streptococcal infections. World Health organization. Geneva; 1996.
4. Davies MR, Tran TN, McMillan DJ, Gardiner DL, Currie BJ, Sriprakash KS. Inter-species genetic movement may blur the epidemiology of streptococcal diseases in endemic regions. *Microbes Infect* 2005;7:1128-38.
5. Menon T, Shanmugasundaram S, Kumar MP, Kumar CP. Group A streptococcal infections of the pharynx in a rural population in south India. *Indian J Med Res* 2004;119:171-3.

Access this article online

Quick Response Code:



Website:

www.jlponline.org

DOI:

10.4103/0974-2727.129099